

On the global generation of direct images of pluri-adjoint line bundles

Masataka Iwai

The University of Tokyo

MSJ Autumn Meeting 2018, at Okayama University.
September 25, 2018

Fujita's freeness conjecture

Conjecture (Fujita's freeness conjecture)

Let X be a smooth projective n -dimensional variety and L be an ample line bundle on X . Is the line bundle $K_X \otimes L^{\otimes n+1}$ globally generated?

Fujita's freeness conjecture

Conjecture (Fujita's freeness conjecture)

Let X be a smooth projective n -dimensional variety and L be an ample line bundle on X . Is the line bundle $K_X \otimes L^{\otimes n+1}$ globally generated?

Fujita's freeness conjecture holds

- if $n = 2$ (Reider 88)
- if $n = 3$ (Ein-Lazarsfeld 93)
- if $n = 4$ (Kawamata 97)
- if $n = 5$ (Ye-Zhu 15)

Fujita's freeness conjecture

Conjecture (Fujita's freeness conjecture)

Let X be a smooth projective n -dimensional variety and L be an ample line bundle on X . Is the line bundle $K_X \otimes L^{\otimes n+1}$ globally generated?

Fujita's freeness conjecture holds

- if $n = 2$ (Reider 88)
- if $n = 3$ (Ein-Lazarsfeld 93)
- if $n = 4$ (Kawamata 97)
- if $n = 5$ (Ye-Zhu 15)

(Angehrn-Siu 95) $K_X \otimes L^{\otimes b}$ is globally generated for any $b \geq \frac{n(n+1)}{2} + 1$.

Conjecture (Popa-Schnell 14)

Let $f: X \rightarrow Y$ be a surjective morphism of smooth projective varieties, with Y of dimension n , and L be an ample line bundle on Y . For any $a \geq 1$, is the sheaf

$$f_*(K_X^{\otimes a}) \otimes L^{\otimes b}$$

globally generated for all $b \geq a(n+1)$?

If $X = Y$ and f is an identity map, Popa-Schnell conjecture implies Fujita's freeness conjecture.

- (Popa-Schnell 14) If L is ample and globally generated, this conjecture holds.

- (Popa-Schnell 14) If L is ample and globally generated, this conjecture holds.
- (Dutta 17) $f_*(K_X^{\otimes a}) \otimes L^{\otimes b}$ is generated by the global sections at a **general point** $y \in Y$ for all $b \geq a\left(\frac{n(n+1)}{2} + 1\right)$.
- (Dutta 17) If f is smooth, $f_*(K_X^{\otimes a}) \otimes L^{\otimes b}$ is globally generated on Y for all $b \geq a\left(\frac{n(n+1)}{2} + 1\right)$.

- (Popa-Schnell 14) If L is ample and globally generated, this conjecture holds.
- (Dutta 17) $f_*(K_X^{\otimes a}) \otimes L^{\otimes b}$ is generated by the global sections at a **general point** $y \in Y$ for all $b \geq a\left(\frac{n(n+1)}{2} + 1\right)$.
- (Dutta 17) If f is smooth, $f_*(K_X^{\otimes a}) \otimes L^{\otimes b}$ is globally generated on Y for all $b \geq a\left(\frac{n(n+1)}{2} + 1\right)$.
- (Deng 17) $f_*(K_X^{\otimes a}) \otimes L^{\otimes b}$ is generated by the global sections at a **general point** $y \in Y$ for all $b \geq n^2 - n + a(n + 1)$.

- (Popa-Schnell 14) If L is ample and globally generated, this conjecture holds.
- (Dutta 17) $f_*(K_X^{\otimes a}) \otimes L^{\otimes b}$ is generated by the global sections at a **general point** $y \in Y$ for all $b \geq a\left(\frac{n(n+1)}{2} + 1\right)$.
- (Dutta 17) If f is smooth, $f_*(K_X^{\otimes a}) \otimes L^{\otimes b}$ is globally generated on Y for all $b \geq a\left(\frac{n(n+1)}{2} + 1\right)$.
- (Deng 17) $f_*(K_X^{\otimes a}) \otimes L^{\otimes b}$ is generated by the global sections at a **general point** $y \in Y$ for all $b \geq n^2 - n + a(n + 1)$.

Even if (X, Δ) is a Kawamata log terminal \mathbb{Q} -pair of a normal projective variety and an effective divisor, their results hold, that is, $f_*\left(\mathcal{O}_X(a(K_X + \Delta))\right) \otimes L^{\otimes b}$ is generated by the global sections at a **general point** $y \in Y$.

Theorem (I.17)

Let $f: X \rightarrow Y$ be a surjective morphism of smooth projective varieties, with Y of dimension n , and L be an ample line bundle on Y .

If y is a **regular value** of f , then for any $a \geq 1$ the sheaf

$$f_*(K_X^{\otimes a}) \otimes L^{\otimes b}$$

is generated by the global sections at y for all $b \geq \frac{n(n-1)}{2} + a(n+1)$.

Theorem (I.17)

Let $f: X \rightarrow Y$ be a surjective morphism of smooth projective varieties, with Y of dimension n , and L be an ample line bundle on Y .

If y is a **regular value** of f , then for any $a \geq 1$ the sheaf

$$f_*(K_X^{\otimes a}) \otimes L^{\otimes b}$$

is generated by the global sections at y for all $b \geq \frac{n(n-1)}{2} + a(n+1)$.

By using result, if f is smooth, $f_*(K_X^{\otimes a}) \otimes L^{\otimes b}$ is globally generated on Y for all $b \geq \frac{n(n-1)}{2} + a(n+1)$.

Theorem (I.17, Dutta-Murayama 17)

Let (X, Δ) be a Kawamata log terminal \mathbb{Q} -pair of a normal projective variety and an effective divisor, and Y be a smooth projective n -dimensional variety.

Let $f: X \rightarrow Y$ be a surjective morphism, and L be an ample line bundle on Y .

For any $a \geq 1$ such that $a(K_X + \Delta)$ is an integral Cartier divisor, the sheaf

$$f_*\left(\mathcal{O}_X(a(K_X + \Delta))\right) \otimes L^{\otimes b}$$

is generated by the global sections at a **general point** $y \in Y$ for all $b \geq \frac{n(n-1)}{2} + a(n+1)$

Even if X is a normal analytic variety in Fujiki's class C , the above theorem holds.

Idea of proof

We show that for any regular value $y \in Y$, any section

$$s \in H^0(X_y, K_{X_y}^{\otimes a} \otimes f^*(L)^{\otimes b}|_{X_y})$$

can be extended to X .

We show that for any regular value $y \in Y$, any section

$$s \in H^0(X_y, K_{X_y}^{\otimes a} \otimes f^*(L)^{\otimes b}|_{X_y})$$

can be extended to X .

- 1 We take a good singular Hermitian metric into $K_X^{\otimes a-1} \otimes f^*(L^{\otimes b})$.

We show that for any regular value $y \in Y$, any section

$$s \in H^0(X_y, K_{X_y}^{\otimes a} \otimes f^*(L)^{\otimes b}|_{X_y})$$

can be extended to X .

- 1 We take a good singular Hermitian metric into $K_X^{\otimes a-1} \otimes f^*(L^{\otimes b})$.
- 2 There exists a neighborhood $V \ni y$ and a section S_V on $f^{-1}(V)$ such that $S_V|_{X_y} = s$. (by using relative version of Ohsawa-Takegoshi L^2 extension.)

Idea of proof

We show that for any regular value $y \in Y$, any section

$$s \in H^0(X_y, K_{X_y}^{\otimes a} \otimes f^*(L)^{\otimes b}|_{X_y})$$

can be extended to X .

- 1 We take a good singular Hermitian metric into $K_X^{\otimes a-1} \otimes f^*(L^{\otimes b})$.
- 2 There exists a neighborhood $V \ni y$ and a section S_V on $f^{-1}(V)$ such that $S_V|_{X_y} = s$. (by using relative version of Ohsawa-Takegoshi L^2 extension.)
- 3 We solve a $\bar{\partial}$ -equation to extend the section S_V .